Perinatal Depression Resource Guide













Table of Contents

Screening for Depression	3	
Risk Factors Checklist	4	
In-Network Experts	5-7	
Why it's important	8	
Resources: During Pregnancy	9	
Resources: After the Baby is Born	10	
Baby Blues v. Postpartum Depression		



Screening for Perinatal Depression



"The vast majority of postpartum women with depression are not identified or treated even though they are at higher risk for psychiatric disorders... A lot of women do not understand what is happening to them. They think they're just stressed or they believe it is how having a baby is supposed to feel"

-Dr. Kathrerine Wisner

-Ur. Kathrerine Wisner
Director of Northwestern University's Asher Centre
for Research and Treatment of Depression

Screening Tool	Number of Items	Time to Complete (Minutes)	Sensitivity and Specificity	Spanish Available
Edinburgh Postnatal Depression Scale*	10	Less than 5	Sensitivity 59-100% Specificity 49-100%	Yes
Postpartum Depression Screening Scale*	35	5-10	Sensitivity 91–94% Specificity 72–98%	Yes
Patient Health Questionnaire 9*	9	Less than 5	Sensitivity 75% Specificity 90%	Yes
Beck Depression Inventory	21	5-10	Sensitivity 47.6-82% Specificity 85.9-89%	Yes
Beck Depression Inventory-II	21	5-10	Sensitivity 56–57% Specificity 97–100%	Yes
Center for Epidemiologic Studies Depression Scale*	20	5-10	Sensitivity 60% Specificity 92%	Yes
Zung Self-rating Depression Scale	20	5-10	Sensitivity 45-89% Specificity 77-88%	No

Souce: http://www.acog.org/Resources-And-Publications/Committee-Opinions/Committee-on-Obstetric-Practice/Screening-for-Perinatal-Depression

*Go to: www.mededppd.org/screening_tools.asp to download

Risk Factors Checklist

□ Being under the age of 20
□ Alcohol, drug, or tobacco use
□ Having an unplanned pregnancy
□ Having mixed feelings about the pregnancy
□ A history of depression, bipolar disorder, or anxiety disorders
□ Having a close family member with depression
□ Experiencing a stressful event during the pregnancy or delivery
□ A bad relationship with your significant other
□ Being single while pregnant
□ Money or housing problems
□ Poor family supports

If you have checked *any* of the above boxes, you are at risk for developing Perinatal Depression. You are not alone; approximately 800,000 women are diagnosed with Perinatal Depression each year.

In-Network Experts

Each provider listed below has a staff member who has participated in the Perinatal Depression Training
Institute and has completed the 6-12 month specialized training course.

They are all considered experts in Perinatal Depression in Philadelphia.



<u>Einstein Medical Center</u> 5501 Old York Road – 215-456-7890

Hispanic Community Counseling 3156 Kensington Avenue #3 - 215-291-8151

> Intercultural Family Services, Inc. 4225 Chestnut Street – 215-386-1298

<u>Progressions Behavioral Health</u> 330 Henry Avenue #302 – 215-924-0684

Stop & Surrender Program 2522 W Huntingdon Street – 215-225-4626

Women's Institute for Family Health

5936 Chestnut Street – 215-747-2391

Stephen and Sandra Sheller 11th Street Family Health Services (RHD & Drexel)

850 N 11th Street - 267-769-1100

Asociación Puertorriqueños en Marcha (APM)

1900 N 9th Street - 267-296-7200

Esperanza Health Centers

2940 N 5th Street - 215-221-6633

3156 Kensington Avenue - 215-831-1100

4417 6th Street - 215-302-3600

Warren E. Smith Health System

1315 Windrim Avenue - 215-455-3900

Thomas Jefferson - Family Center

1233 Locust Street - 215-955-8577

COMHAR

100 W Lehigh Avenue - 215-203-3000

Best Behavioral Healthcare

5043 Frankford Avenue - 215-744-4343

Consortium

5501 Chestnut Street - 215-596-8100

Pan American

2561 N Front Street - 215-425-8909

City Health Centers

Health Center 3 555 S. 43rd Street – 215-685-7504

Health Center 4 4400 Haverford Avenue – 215-685-7601

> Health Center 5 1900 N 20th Street – 215-685-2933

Health Center 6 301 West Girard Avenue – 215-685-3808


Health Center 9 131 E. Chelten Avenue – 215-685-5701

Health Center 10 2230 Cottman Avenue – 215-685-0639

Why It's Important:

Facts about Perinatal Depression

National Survey of Drug Use and Health (NSDUH) data show that "women suffering from depression are at increased risk of substance abuse, developing chronic diseases, and having poorer health" (Creanga et. Al. 2014)



"Suicide accounted for about 20 percent of postpartum deaths and is the second most common cause of mortality in postpartum women" -www.upi.com

Resources: During Pregnancy

National Child & Maternal Health Education Program: https://www.nichd.nih.gov/ncmhep/MMHM/Pages/index.aspx

2020 Mom Project: http://www.2020mom.org/



Centers for Disease Control and Prevention: http://www.cdc.gov/reproductivehealth/MaternalInfantHealth/PMSS.html

Maternity Care Coalition: www.MaternityCareCoalition.org
Cribs for Kids, Early Head Start, Healthy Families America, Healthy Start, HMO Outreach
Program, MoMobile, NPhila Breastfeeding Program, Parenting Collaborative Groups, Safe Start for Moms

Resources: After the Baby is Born

Postpartum Support International: http://www.postpartum.net/

Postpartum Progress: http://www.postpartumprogress.com/

The Benefit Bank: <u>www.thebenefitbank.com</u> online resource to help families link to public benefits

Parent University: www.philasd.org/parentuniversity classes and workshops for parents through the School District

Community Legal Services: www.cls.phila.org

Free legal services

Greater Philadelphia Coalition Against Hunger: www.hungercoalition.org
Emergency food resources and other food programs

Smoke Free Philly: 215-683-LIVE

Text for Baby: www.text4baby.org

Free weekly text messages to help mothers through pregnancy and through baby's first year. Text

BABY to 511411 or BEBE for Spanish

Take Control Philly: www.takecontrolphilly.org

Information on STDs and safe sex



Baby Blues	OR	Postpartum Depression
Weepiness/crying for no apparent reason		Difficulty sleeping, insomnia, exhaustion
Feeling "I'm not like myself" or "this isn't me"		Lack of energy
Impatience		Low sex drive
Irritability		Changes in appetite, weight loss, weight gain
Restlessness		Weepiness, excessive worry, agitation, anxiety
Anxiety		Feelings of inadequacy, hopelessness, despair
Sadness		Guilt, sadness, fear of being alone
Mood changes		Irritability
Poor concentration		Difficulty concentrating, panic, anger
		Scary thoughts about baby, over-concern for baby's health